

Available online at www.sciencedirect.com



DOMESTIC ANIMAL ENDOCRINOLOGY

Domestic Animal Endocrinology 29 (2005) 23-33

www.journals.elsevierhealth.com/periodicals/dae

The anterior pituitary gland: Lessons from livestock

C.G. Scanes ^{a,b,*}, S. Jeftinija ^c, A. Glavaski-Joksimovic ^d, J. Proudman ^e, C. Arámburo ^f, L.L. Anderson ^{b,d}

- a Department of Poultry Science, Mississippi State University, 617 Allen Hall, P.O. Box 6343, MS 39762, USA
- b Department of Basic Science, Mississippi State University, MS 39762, USA
 c Department of Animal Science, Department of Biomedical Sciences,
 Iowa State University, Ames, IA 50011, USA
- ^d Department of Biomedical Sciences, Iowa State University, Ames, IA 50011, USA
 ^e Biotechnology and Germplasm Laboratory, Agricultural Research Service,
 US Department of Agriculture, Beltsville, MD 20705, USA
- f Department of Ĉellular and Molecular Neurobiology, Institute of Neurobiology, Universidad Nacional Autónoma de México, Campus UNAM-UAQ, Juriquilla, Querétaro, México

Abstract

There has been extensive research of the anterior pituitary gland of livestock and poultry due to the economic (agricultural) importance of physiological processes controlled by it including reproduction, growth, lactation and stress. Moreover, farm animals can be biomedical models or useful in evolutionary/ecological research. There are for multiple sites of control of the secretion of anterior pituitary hormones. These include the potential for independent control of proliferation, differentiation, de-differentiation and/or inter-conversion cell death, expression and translation, post-translational modification (potentially generating multiple isoforms with potentially different biological activities), release with or without a specific binding protein and intra-cellular catabolism (proteolysis) of pituitary hormones. Multiple hypothalamic hypophysiotropic peptides (which may also be produced peripherally, e.g. ghrelin) influence the secretion of the anterior pituitary hormones. There is also feedback for hormones from the target endocrine glands. These control mechanisms show broadly a consistency across species and life stages; however, there are some marked differences. Examples from growth hormone, prolactin, follicle stimulating hormone and luteinizing

^{*} Corresponding author. Tel.: +1 662 325 3570; fax: +1 662 325 8028. E-mail address: scanes@research.msstate.edu (C.G. Scanes).

hormone will be considered. In addition, attention will be focused on areas that have been neglected including the role of stellate cells, multiple sub-types of the major adenohypophyseal cells, functional zonation within the anterior pituitary and the role of multiple secretagogues for single hormones.

© 2005 Published by Elsevier Inc.

Keywords: Pituitary; LH; GH; Prolactin variants

1. Introduction

This review will consider the structure of the anterior pituitary gland, the control of the secretion of anterior pituitary hormone and the role of isoforms of these hormones. Attention will be focused on areas where livestock and poultry show specific advantages (see Section 5) and also to arenas that have until recently been neglected.

2. Cells in the anterior pituitary gland

The anterior pituitary gland is composed of a mixture of secretory cells together with non-secretory folliculo-stellate cells (FS). In sheep and horses, this profile is observed in both fetal development and post-natal growth [1,2]. In chickens, FS are not detected until post-hatching [3]. It is likely that sub-types of the major cells (corticotrophs, gonadotrophs, lactotrophs, somatotrophs and thyrotrophs together with FS) exist with different functional characteristics. The possibility of there being zonation/anatomical separation of the cell sub-types has received relatively little attention in livestock species. Zonation would facilitate "point to point" control with specific peptides released from areas within the median eminence and passing via designated portal blood vessels to stimulate specific secretory cells.

2.1. Secretory cells

The anterior pituitary gland is "classically" considered to be composed of the following secretory cells: corticotrophs [producing adenocorticotropic hormone (ACTH)]; gonadotrophs [producing luteinizing hormone (LH) and follicle stimulating hormone (FSH)], lactotrophs or mammotroph [producing prolactin]; somatotrophs [producing growth hormone (GH)]; and thyrotrophs [producing thyrotropin (TSH)]. This situation broadly pertains to livestock and poultry but with a number of exceptions, some of which will be addressed.

2.1.1. Gonadotrophs

There is frequently the assumption that gonadotrophs produce both LH and FSH. This is not necessarily the case. In poultry, there are at least two distinct populations of gonadotrophs, producing, respectively, LH and FSH [4] (discussed in more detail by Berghman in this volume). Moreover, these have distinctly different embryonic origens [5]. In sheep, sub-populations of gonadotrophs exit, some containing both LH and FSH and other

only LH [6]. It is not known whether different sub-populations of gonadotrophs produce specific isoforms of LH and/or FSH.

2.1.2. Lactotrophs and somatotrophs

In addition to the somatotrophs and lactotrophs, there is another cell-type in the anterior pituitary gland, the somatolactotroph or somatomammotroph. This synthesizes both GH and prolactin. This cell-type can be an intermediate in the conversion of somatotrophs to lactotrophs for instance in turkeys when starting to exhibit incubation behavior [7,8] or in the opposite direction at the cessation of incubation [9]. It is not clear the extent the release of GH and prolactin can be independently controlled from somatolactotrophs.

The control of the differentiation of somatotrophs has been extensively examined using chick embryo pituitary cells (discussed in detailed by Porter and colleagues elsewhere in this volume). There are marked changes in the number of somatotrophs in the anterior pituitary gland during development of the chick embryo and during post-hatching growth [10–13]. Moreover, the somatotrophs undergo changes with increases in secretory granules [13]. The percentage of reaches a plateau of ~40% of the cells in the caudal lobe of the anterior pituitary gland which is maintained despite the continued growth of the anterior pituitary gland [13]. In adult turkeys, there are increases in the number and proliferation of somatotrophs following the termination of incubating behavior when the number of lactotroph is declining [9]. These studies would suggest a "set point" for the number/proportion of somatotrophs and a mechanism(s) controlling somatotroph number/percentage. Moreover, it would also provide support for GH having importance in late growth and in the adult.

It may be questioned whether different sub-types of somatotrophs exist and are stimulated by different secretagogues. Multiple hypothalamic peptides stimulate GH release from the somatotroph. These include the following: GH releasing hormone (GHRH), ghrelin and pituitary adenylate cyclase activating peptide (PACAP) acting, respectively, via GHRH receptors, the GH secretagogue (GHS) receptor (GHSR) and PACAP receptor and increasing intra-cellular calcium (see Table 1) (e.g. [14]). In addition, other peptides increase intra-cellular calcium in somatotrophs. These include leptin (pigs and chickens), motilin (pigs) and thyrotropin releasing hormone (chicken but not pigs). What is interesting is that not all somatotrophs respond to all secretagogues (see Table 1). This might support that the existence of different sub-types of somatotrophs responding

Table 1
Percentage of porcine and chicken somatotropes (defined based on ability of GHRH to elicit an increase in the intra-cellular calcium concentration) responding to various other secretagogues by increase in the intra-cellular calcium concentration

	Porcine	Chicken
GHRH, 10 μM	100 ^a	100a
PACAP, 1 μM	96	85
TRH, 1 μM	N.A.	73
Ghrelin, 1 μM	98	12
Leptin	54	58

^a By definition.

to different hypothalamic secretagogues. It is possible that different isoforms of GH or prolactin are produced by various sub-populations of somatotrophs, lactotrophs and/or somatolactotrophs.

2.2. Folliculo-stellate cells (FS)

There is increasing interest in the FS exerting a paracrine control of secretory cells in the anterior pituitary gland. However, this has not received much attention in livestock species as yet. The presence of G-100 protein is frequently used to distinguish FS. Glial fibrillary acidic protein is a marker for astrocytes. However, this protein is not detected in FS in the anterior pituitary gland in some studies (e.g. cattle [15]) but is in others (rat [16]). The FS exert a paracrine effect on the secretory cells based on rodent studies (reviewed [17]). For example, growth of lactotrophs can stimulated by indirectly by transforming growth factor-beta3 acting on FS which in turn release fibroblast growth factor [18]. FS also influence hormone release from secretory cells; mediating effects of steroids on prolactin release (reviewed [19]). Similarly, interferon-gamma inhibits prolactin release with the effect mediated by FS [20]. In addition, FS produce interleukin 6 in response to calcitonin [21]. There is little work on the role of FS in pituitary functioning in livestock species except that bovine FS have been shown to have β-adrenergic receptors coupled to adenylate cyclase [22] and to produce vascular endothelial growth factor and follistatin (Gospodarowicz and Lau [23]).

There is presently untapped potential to use of the pituitary FS in livestock for biomedical models. For instance, the FS of cattle have been employed as models for the proteome of FS with metallothionein I-II immunoreactivity detected recently [24]. Moreover, there appears to be a link between FS and aging based on studies in the pig. During aging, there are marked increases in colloid filled follicles in the anterior pituitary gland [25,26]. The FS, containing G-100 protein, surround the colloid filled follicles and are located in close association with lactotrophs and gonadotrophs [25,26]. It is reasonable, therefore, to propose the pig pituitary and its FS as a model for endocrine aging. Moreover, poultry models may be useful in determining the role of FS. In the chicken, folliculo-stellate cells, as detected by S-100 immunoreactivity, have been reported in the both the cephalic and caudal lobes of the anterior pituitary gland [27]. The FS are in contact with secretory cells that produce LH, GH, POMC and prolactin [1]. There are changes in FS with physiological state. For instance, FS are not detected in the pituitary of the chick embryo but are apparent by 4 weeks of age [1]. Moreover, there are changes in the FS in the caudal lobe of the anterior pituitary following feed withdrawal induction of molt in the hen; the amount of G-100 immunorectivity (area of staining) increasing initially following resumption of lay [27]. Moreover, challenges with Escherichia coli have been associated with necrosis of FS in the anterior pituitary gland [28].

3. Control of secretion of anterior pituitary hormones

There is the potential for multiple sites of control of the secretion of anterior pituitary hormones. These include the potential for independent control of the following: (1) proliferation, differentiation, de-differentiation and/or inter-conversion, cell death of specific populations of secretory cells; (2) expression and translation; (3) post-translational

modification (potentially generating multiple isoforms); (4) release with or without a specific binding protein and (5) intra-cellular catabolism (proteolysis) of pituitary hormones.

Geoffrey Harris conceived the physiological system by which the hypothalamus controls the release of hormones from the anterior pituitary gland with releasing factors from the median eminence passing through the hypophyseal portal blood vessels to the secretory cells (reviewed [29]). The concept was essentially that a single releasing factor or hormone acts in an endocrine manner to elicit release of a hormone with the response proportional to the dose. It is increasingly clear that the is secretion of anterior pituitary hormones under control of the following mechanisms:

- 1. Multiple hypothalamic releasing hormones that may act on secretory cells in specific zones of the anterior pituitary gland;
- 2. Paracrine factors within the anterior pituitary gland including the putative influence of the FS (see discussion of FS and excitory amino-acids);
- 3. Peripheral hormones/factors acting directly on the secretory cells or FS.

The latter includes hormones from target tissues that exert feedback effects both at the level of the hypothalamus and anterior pituitary gland (negative feedback: sex steroids on LH and FSH release, inhibin on FSH release, triiodothyronine on TSH release, insulin-like growth factor-1 on GH release and glucocorticoids on ACTH release). In addition, hormone secretion from secretory cells is stimulated by positive feedback acting at the level of the hypothalamus (e.g. in female mammals, estradiol and, in poultry, progesterone stimulating the pre-ovulatory LH surge). In addition to the feedback effects, there are peptides that are produced by peripheral tissues and influence hormone secretion from the anterior pituitary gland. These may or may not be present in the hypothalamus and/or released into the portal blood vessels. Examples of such peptides that influence the release of GH include somatostatin produced by the pancreas and gastro-intestinal tract, ghrelin produced by the stomach, GHRH produced by the placenta and leptin produced by the adipose tissue and liver.

3.1. Peptides

While initially there was the assumption none releasing factor for each adenohyophyseal hormone, the situation is becoming increasingly complex. The release of LH, TSH and ACTH are, respectively, stimulated by gonadotropin releasing hormone (GnRH), thyrotropin releasing hormone (TRH) and corticotropin releasing hormone (CRH). The situation with FSH is problematic. The release of both LH and FSH is stimulated by GnRH in mammals (reviewed see [30,67]). The lamprey GnRH III (lGnRh III) has been proposed as the endogenous specific FSH releasing factor ([31,32], also reviewed [33]) and appears to act in that way in cattle [34]. There are three decapeptide releasing hormones for LH found in the avian hypothalamus; namely cGnRH I, cGnRH II and lGnRh III [35–37,8]. We recently examine compared the ability of the three GnRH peptides to evoke release of LH and FSH in adult male chickens (broiler breeders). Markedly increased circulating concentrations of LH were observed with challenges with cGnRH I or II. These peptides also increased the circulating concentration of FSH but the response was much less robust.

Isoform	Immunological (RIA) potency ^a	Biological activity		
		Progesterone synthesis in vitro ^{a, b}	Ovulation induction in vivo ^c	
			Low dose	High dose
1	220	12.3	0/2 (30 μg)	2/2 (116 μg)
2	137	8.0	$0/2 (34 \mu g)$	$1/2 (67 \mu g)$
3	137	4.1	$0/2 (49 \mu g)$	$2/2 (96 \mu g)$
4	110	4.4	$0/2 (36 \mu g)$	2/2 (72 μg)
5	110	1.8	$1/2 (15 \mu g)$	$2/2 (30 \mu g)$
6	73	N.A.	$0/2 (54 \mu g)$	2/2 (222 μg)

Table 2
Immunological and biological activities of turkey LH isoform (weights adjusted for protein content)

N.A., not available.

In contrast, lGnRH III evoked only a small response in circulating concentrations of LH and no change in those of FSH.

Despite the supposedly straight-forward roles for both GH and prolactin, there are multiple releasing hormones for each. Control for prolactin release is via a series of releasing factors including vasoactive intestinal peptide (VIP)(+), PACAP(+), TRH(+) and dopamine(-). The releasing hormones for GH include GHRH, ghrelin, PACAP and TRH (all stimulatory) and somatostatin (inhibitory) (see Table 2). It might be questioned on what basis do we call the peptide, GHRH, as ghrelin or PACAP also stimulate GH release? Similarly, on what basis do we call TRH as the peptide also stimulates release of PRL (mammals) and GH (birds) and CRH stimulates TSH release in birds?

3.2. Excitatory amino-acids

There is substantial evidence that excitatory amino-acids such as *N*-methyl-D-asparate (NMDA) stimulate the release of anterior pituitary hormones in rodent models. There is, however, a dearth of information on the direct effects of these in livestock. Consider the case of GH, NMDA increases its release from rat pituitary cells (e.g. [38,39,11,40–42]). Similarly, in vivo studies in livestock indicate that NMDA increases GH secretion (cattle [43]; pigs [44,45]; horses [46]). It is not known whether these effects in livestock reflect direct effects on the somatotrophs or indirect effects via the hypothalamus. There is, however, no information in the direct effects of NMDA on somatotrophs of any livestock species. Moreover, there are no reports of the effects of NMDA influencing the release of pituitary hormone in poultry except that NMDA has been observed to influence the release of GnRH from hypothalamic tissue in vitro [47]. It is not clear whether the excitatory amino-acids that influence GH release originate in the hypothalamus and/or cells within the anterior pituitary gland, for instance, the FS.

^a Potency is expressed as a percentage relative to a turkey LH standard at 50% B/B_0 for RIA and ED₅₀ for bioassay (B221B).

^b Assayed based on the in vitro chicken granulosa cell assay using progesterone production as the end point [2] (F. Hertelendy and J. Proudman, unpublished observations).

^c Data is expressed as the number of turkey hens exhibiting premature ovulation in response to stimulus. Ovine LH failed to induce ovulation at 500, 1000 and 2000 µg doses (data from H. Opel and J. Proudman, unpublished).

4. Isoforms to proteomics

There has been tremendous progress made in our understanding of gene expression in the anterior pituitary gland [48]. In the near future, there will be considerable attention focused on the pituitary proteome, particularly at the individual cell level. A substantial number of biologically active peptides have been and it is predicted will be found in the anterior pituitary gland. For instance, pancreastatin has been found in the pig pituitary gland [12]. In contrast to the strong interest in the biomedical significance of post-translational isoforms of anterior pituitary hormones, these isoforms in livestock and poultry have received less attention. Post-translational isoforms of LH exist with, for instance chicken and porcine LH [49,13]. These have different biological activities (for chicken LH, see Table 2). Similarly, there is evidence of different isoforms of FSH with different biological activities that are released in livestock mammals [50,51]; also see review [52] and of the α sub-unit of LH, FSH and TSH [53]. There are multiple post-translational isoforms of prolactin. These include the following: glycosylated (e.g. ovine [54], porcine [55], turkey [56]), phosphorylated (e.g. bovine [68]; cleaved prolactin [69] and cleaved and reduced fragments of prolactin (e.g. rat 16 kDa [57]) These isoforms can have different biological activities. For instance, glycosylated prolactin has reduced pigeon crop assay activity [58]. Phosphorylated prolactin exhibits different activities than non-phosphorylated prolactin, not being luteotropic but also antagonizing the mitogenic effects of prolactin [59–61]. The 16 kDa rat prolactin has a low prolactin activity [57] but exhibits anti-angiogenic activity ([62]; reviewed [63]). The relative concentrations of different isoforms can exhibit marked changes with physiological state (e.g. glycosylated versus non-glycosylated prolactin in pigs [64,65] and turkeys [6,7]).

A similar situation exists for GH. Using the example of the chicken, GH isoforms include the following: (1) a splicing variant expressed in the eye [66] which analogous to the 20 kDa human GH and (2) post-translationally modified isoforms, e.g. phosphorylated [3,4], glycosylated [9], dimeric and other oligomeric forms [4], proteolytically cleaved and both cleaved and reduced to yield a 15 kDa fragment [5]. Monomeric GH binds to the GH receptor [4,70] and is biologically active—stimulating hepatic IGF-1 release and lipolysis [70]. The glycosylated GH is biologically active but has a different clearance rate [10]. No GH receptor binding activity is seen with dimer GH, other oligomers of GH or the 15 kDa fragment of GH [4,5]. However, the 15 kDa GH isomer/fragment has angiogenic activity as indicated by stimulation of proliferation of endothelial cells [5]. It might be noted that the profile of GH isoforms in the pituitary exhibit changes during growth and development with more 15 kDa in the embryonic pituitary gland [4].

5. Advantages of domestic animals in pituitary research

There are advantages of using livestock and poultry for endocrine research. These include the following: (1) the agricultural/economic importance and hence potential funding (government and private industry) of the reproduction, growth, milk production (mammogenesis, lactation, etc.) and responses to stress; (2) availability of disparate genetic breeds, varieties and lines with known genetics (including polymorphisms in candidate genes, e.g. SNPS); (3) the established body of knowledge on interactions between genotype, phenotype and

environment; and (4) that livestock and poultry represent multiple evolutionarily lines separated by about 200 million years ago. Disadvantages of using livestock include the lack of "knock-out models" and the difficulty in maintaining livestock in many biomedical facilities.

References

- Allaerts W, Boonstra-Blom AG, Peeters K, Janse EM, Berghman LR, Jeurissen SH. Prenatal development of hematopoietic and hormone-producing cells in the chicken adenohypophysis. Gen Comp Endocrinol 1999;114:213–24.
- [2] Aramburo C, Donoghue D, Montiel JL, Berghman IR, Scanes CG. Phosphorylation of chicken growth hormone. Life Sci 1990;47:945–52.
- [3] Arámburo C, Montiel JL, Proudman JA, Berghman LR, Scanes CG. Phosphorylation of PRL and growth hormone. J Mol Endocrinol 1992;8:181–91.
- [4] Arámburo C, Luna M, Carranza M, Reyes M, Martinez-Coria H, Scanes CG. Growth hormone size variants: changes in the pituitary during development of the chicken. Proc Soc Exp Biol Med 2000;223:67–74.
- [5] Arámburo C, Carranza M, Reyes M, Luna M, Coria H, Berúmen L, et al. Characterization of a bioactive 15 kDa fragment produced by proteolytic cleavage of chicken growth hormone. Endocrine 2001;15:231–40.
- [6] Bedecarrats G, Guemene D, Morvan C, Crisostomo-Pinto S, Kuhnlein U, Zadworny D. In vitro release of isoforms of PRL from pituitary glands of turkey hens at different physiological stages. Gen Comp Endocrinol 1999;113:105–11.
- [7] Bedecarrats G, Guemene D, Kuhnlein U, Zadworny D. Changes in levels of immunoreactive PRL isoforms during a reproductive cycle in turkey hens. Comp Endocrinol 1999;113:96–104.
- [8] Bentley GE, Moore IT, Sower SA, Wingfield JC. Evidence for a novel gonadotropin-releasing hormone in the hypothalamic and forebrain areas in songbirds. Brain Behav Evol 2004;63:34–46.
- [9] Berghman LR, Lens P, Decuypere E, Kuhn ER, Vandesande F. Glycosylated chicken growth hormone. Gen Comp Endocrinol 1987;68:408–14.
- [10] Berghman LR, Buyse J, Huybrechts LM, Darras VM, Vandesande F, Kuhn ER, et al. Disappearance rate of glycosylated and non-glycosylated chicken growth hormone: influence on biological activity. Comp Biochem Physiol 1994;108C:161–9.
- [11] Bhat GK, Mahesh VB, Chu ZW, Chorich LP, Zamorano PL, Brann DW. Localization of the N-methyl-D-aspartate R1 receptor subunit in specific anterior pituitary hormone cell types of the female rat. Neuroendocrinology 1995;62:178–86.
- [12] Bretherton-Watt D, Ghatei MA, Bishop AE, Facer P, Fahey M, Hedges M, et al. Pancreastatin distribution and plasma levels in the pig. Peptides 1988;9:1005–14.
- [13] Burova T, Lecompte F, Galet C, Monsallier F, Delpech S, Haertle T, et al. Conformational stability and in vitro bioactivity of porcine luteinizing hormone. Mol Cell Endocrinol 2001;176:129–34.
- [14] Glavaski-Joksimovic A, Jeftinija K, Scanes CG, Anderson LL, Jeftinija S. Stimulatory effect of ghrelin on isolaterd porcine somatotropes. Neuroendocrinology 2003;77:366–78.
- [15] Shimada T. Immunohistochemical localization of keratin in bull, goat, and sheep anterior pituitary glands. Cell Tissue Res 1992;267:251–60.
- [16] Van Nassauw L, Harrisson F, Cras P, Callebaut M. Immunohistochemical localization of S-100 protein, glial fibrillary acidic protein, and neuron-specific enolase in the pars distalis of quail, rat, and human hypophyses. Histochemistry 1987;86:353–8.
- [17] Vankelecom H, Denef C. Paracrine communication in the anterior pituitary as studied in reaggregate cell cultures. Microsc Res Tech 1997;39:150–6.
- [18] Hentges S, Boyadjieva N, Sarkar DK. Transforming growth factor-beta3 stimulates lactotrope cell growth by increasing basic fibroblast growth factor from folliculo-stellate cells. Endocrinology 2000;141:859–67.
- [19] Morris JF, Christian HC, Chapman LP, Epton MJ, Buckingham JC, Ozawa H, et al. Steroid effects on secretion from subsets of lactotrophs: role of folliculo-stellate cells and annexin 1. Physiol Biochem 2002;110:54–61.
- [20] Vankelecom H, Andries M, Billiau A, Denef C. Evidence that folliculo-stellate cells mediate the inhibitory effect of interferon-gamma on hormone secretion in rat anterior pituitary cell cultures. Endocrinology 1992;130:3537–46.

- [21] Kiriyama Y, Tsuchiya H, Murakami T, Satoh K, Tokumitsu Y. Calcitonin induces IL-6 production via both PKA and PKC pathways in the pituitary folliculo-stellate cell line. Endocrinology 2001;142:3563–9.
- [22] Findell PR, Weiner RI. Bovine pituitary folliculo-stellate cells have beta-adrenergic receptors positively coupled to adenosine 3',5'-cyclic monophosphate production. Endocrinology 1988;123:2454–61.
- [23] Gospodarowicz D, Lau K. Pituitary follicular cells secrete both vascular endothelial growth factor and follistatin. Biochem Biophys Res Commun 1989;165:288–92.
- [24] Zatt P, Zambenedetti P, Wittkowski W, Carpene E. Localization of metallothionein I–II immunoreactivity in bovine pituitary gland. Life Sci 2001;70:659–67.
- [25] Ogawa S, Couch EF, Kubo M, Sakai T, Inoue K. Histochemical study of follicles in the senescent porcine pituitary gland. Arch Histol Cytol 1996;59:467–78.
- [26] Ogawa S, Ishibashi Y, Sakamoto Y, Kitamura K, Kubo M, Sakai T, et al. The glycoproteins that occur in the colloids of senescent porcine pituitary glands are clusterin and glycosylated albumin fragments. Biochem Biophys Res Commun 1997;234:712–8.
- [27] Chowdhury VS, Yoshimura Y. Changes in the population of immunoreactive S-100-positive folliculo-stellate cells in hens during induced molting. Poult Sci 2002;81:556–60.
- [28] Fernandez AJ, Sierra MA, Mendez A, Mozos E, Moyano MC, Carrasco L. Ultrastructural modifications of the cavities formed by folliculo-stellate cells in chicken adenohypophysis under septic shock conditions. Cell Struct Funct 1986;11:3782–93.
- [29] Raisman G. An urge to explain the incomprehensible: Geoffrey Harris and the discovery of the neural control of the pituitary gland. Annu Rev Neurosci 1997;20:533–66.
- [30] McNeilly AS, Crawford JL, Taragnat C, Nicol L, McNeilly JR. The differential secretion of FSH and LH: regulation through genes, feedback and packaging. Reprod Suppl 2003;61:463–76.
- [31] Yu WH, Karanth S, Walczewska A, Sower SA, McCann SM. A hypothalamic follicle-stimulating hormonereleasing decapeptide in the rat. Proc Natl Acad Sci USA 1997;94:9499–503.
- [32] Yu WH, Karanth S, Sower SA, Parlow AF, McCann SM. The similarity of FSH-releasing factor to lamprey gonadotropin-releasing hormone III (I-GnRH-III). Proc Soc Exp Biol Med 2000;224:87–92.
- [33] McCann SM, Karanth S, Mastronardi CA, Dees WL, Childs G, Miller B, et al. Hypothalamic control of gonadotropin secretion. Prog Brain Res 2002;141:151–64.
- [34] Dees WL, Dearth RK, Hooper RN, Brinsko SP, Romano JE, Rahe H, et al. Lamprey gonadotropin-releasing hormone-III selectively releases follicle stimulating hormone in the bovine. Domest Anim Endocrinol 2001;20:279–88.
- [35] King JA, Millar RP. Structure of chicken hypothalamic luteinizing hormone-releasing hormone. I. Structural determination on partially purified material. J Biol Chem 1982;257:10722–8.
- [36] King JA, Millar RP. Structure of chicken hypothalamic luteinizing hormone-releasing hormone. II. Isolation and characterization. J Biol Chem 1982;257:10729–32.
- [37] Millar RP, Milton RC, Follett BK, King JA. Receptor binding and gonadotropin-releasing activity of a novel chicken gonadotropin-releasing hormone ([His5, Trp7, Tyr8]GnRH) and a D-Arg6 analog. Endocrinology 1986;119:224–31.
- [38] Lindstrom P, Ohlsson L. Effect of N-methyl-D,L-aspartate on isolated rat somatotrophs. Endocrinology 1992;131:1903–7.
- [39] Niimi M, Sato M, Murao K, Takahara J, Kawanishi K. Effect of excitatory amino acid receptor agonists on secretion of growth hormone as assessed by the reverse haemolytic plaque assay. Neuroendocrinology 1994;60:173–8.
- [40] Pinilla L, Gonzalez L, Tena-Sempere M, Dieguez C, Aguilar E. Gonadal and age-related influences on NMDA-induced growth hormone secretion in male rats. Neuroendocrinology 1999;69:11–9.
- [41] Pinilla L, Gonzalez LC, Tena-Sempere M, Aguilar E. Cross-talk between excitatory and inhibitory amino acids in the regulation of growth hormone secretion in neonatal rats. Neuroendocrinology 2001;73:62–7.
- [42] Pinilla L, Barreiro ML, Tena-Sempere M, Aguilar E. Role of ghrelin in the control of growth hormone secretion in prepubertal rats: interactions with excitatory amino acids. Neuroendocrinology 2003;77:83– 90.
- [43] Shahab M, Nusser KD, Griel LC, Deaver DR. Effect of a single intravenous injection of *N*-methyl-D,L-aspartic acid on secretion of luteinizing hormone and growth hormone in Holstein bull calves. J Neuroendocrinol 1993;5:469–73.

- [44] Estienne MJ, Hurlock WF, Barb CR. Serum concentrations of luteinizing hormone, growth hormone, and cortisol in gilts treated with N-methyl-D,L-aspartate during the estrous cycle or after ovariectomy. J Anim Sci 1998;76:2162–8.
- [45] Estienne MJ, Barb CR. Modulation of growth hormone, luteinizing hormone, and testosterone secretion by excitatory amino acids in boars. Reprod Biol 2002;2:13–24.
- [46] Sticker LS, Thompson Jr DL, Gentry LR. Pituitary hormone and insulin responses to infusion of amino acids and N-methyl-D,L-aspartate in horses. J Anim Sci 2001;79:735–44.
- [47] Jacquet JM, Robinson A, Caldani M. Release of chicken luteinising hormone-releasing hormone-I (cLHRH-I) by mediobasal hypothalamus in the cockerel: validation of an incubation system and effect of excitatory amino acids. Reprod Nutr Dev 1994;34:185–91.
- [48] Cogburn LA, Wang X, Carre W, Rejto L, Aggrey SE, Duclos M, et al. Functional genomics in chickens: development of integrated-systems microarrays for transcriptional profiling and discovery of regulatory pathways. Comp Funct Genomics 2004;5:253–61.
- [49] Krishnan KA, Proudman JA, Bahr JM. Purification and partial characterization of isoforms of luteinizing hormone from the chicken pituitary gland. Comp Biochem Physiol Biochem Mol Biol 1994;108:253– 64.
- [50] Cooke DJ, Crowe MA, Roche JF. Circulating FSH isoform patterns during recurrent increases in FSH throughout the oestrous cycle of heifers. J Reprod Fertil 1997;110:339–45.
- [51] West CR, Carlson NE, Lee JS, McNeilly AS, Sharma TP, Ye W, et al. Acidic mix of FSH isoforms are better facilitators of ovarian follicular maturation and E2 production than the less acidic. Endocrinology 2002;143:107–16.
- [52] Padmanabhan V, Lee JS, Beitins IZ. Follicle-stimulating isohormones: regulation and biological significance. J Reprod Fertil Suppl 1999;54:87–99.
- [53] Chabot V, Magallon T, Taragnat C, Combarnous Y. Two free isoforms of ovine glycoprotein hormone alphasubunit strongly differ in their ability to stimulate prolactin release from foetal pituitaries. J Endocrinol 2000;164:287–97.
- [54] Lewis UJ, Singh RMP, Lewis LJ, Seavey BK, Sinha YN. Glycosylated ovine PRL. Proc Natl Acad Sci 1984;81:385–9.
- [55] Sinha YN, DePaolo LV, Haro LS, Singh RN, Jacobsen BP, Scott KE, et al. Isolation and biochemical properties of four forms of glycosylated porcine PRL. Mol Cell Endocrinol 1991;80:203–13.
- [56] Corcoran DH, Proudman JA. Isoforms of turkey PRL: evidence for differences in glycosylation and in tryptic peptide mapping. Comp Biochem Physiol B 1991;99:563–70.
- [57] Clapp C, Sears PS, Nicoll CS. Binding studies with intact rat PRL and a 16K fragment of the hormone. Endocrinology 1989;125:1054–9.
- [58] Lewis UJ, Singh RN, Lewis LJ. Two forms of glycosylated human PRL have different pigeon crop sacstimulating activities. Endocrinology 1989;124:1558–63.
- [59] Wang YF, Walker AM. Dephosphorylation of standard PRL produces a more biologically active molecular. Evidence for antagonism between non-phosphorylated and phosphorylated PRL in the stimulation of Nb2 cell proliferation. Endocrinology 1993;133:2156–60.
- [60] Yang L, Kuo CB, Liu Y, Coss D, Xu X, Chen C, et al. Administration of unmodified PRL (U-PRL) and a molecular mimic of phosphorylated PRL (PP-PRL) during rat pregnancy provides evidence that the U-PRL:PP-PRL ratio is crucial to the normal development of pup tissues. J Endocrinol 2001;168:227–38.
- [61] Xu X, Wu W, Williams V, Khong A, Chen YH, Deng C, et al. Opposite effects of unmodified PRL and a molecular mimic of phosphorylated PRL on morphology and the expression of prostate specific genes in the normal rat prostate. Prostate 2003;54:25–33.
- [62] Ferrara N, Clapp C, Weiner R. The 16K fragment of PRL specifically inhibits basal or fibroblast growth factor stimulated growth of capillary endothelial cells. Endocrinology 1991;129:896–900.
- [63] Corbacho AM, Martinez De La Escalera G, Clapp C. Roles of PRL and related members of the PRL/growth hormone/placental lactogen family in angiogenesis. J Endocrinol 2002;173:219–38.
- [64] Sinha YN, Klemcke HG, Jacobsen BP, Yen JT. Concentrations of glycosylated PRL in the pituitary gland and plasma of lean and obese barrows. Horm Metab Res 1989;21:652–7.
- [65] Sinha YN, Klemcke HG, Maurer RR, Jacobsen BP. Changes in the glycosylated and nonglycosylated forms of PRL and growth hormone in lean and obese pigs during pregnancy. Endocrinology 1990;127:410–8.

- [66] Takeuchi S, Haneda M, Teshigawara K, Takahashi S. Identification of a novel GH isoform: a possible link between GH and melanocortin systems in the developing chicken eye. Endocrinology 2001;142:5158–66.
- [67] Padmanabhan V, Karsch FJ, Lee JS. Hypothalamic, pituitary and gonadal regulation of FSH. Reprod Suppl 2002;59:67–82.
- [68] Brooks CL, Kim BG, Aphale P, Kleeman BE, Johnson GC. Phosphorylated variant of bovine prolactin. Mol Cell Endocrinol 1990;71:117–23.
- [69] Sinha YN, Gilligan TA. A cleaved form of prolactin in the mouse pituitary gland: identification and comparison of in vitro synthesis and release in strains with high and low incidences of mammary tumors. Endocrinology 1984;114:2046–53.
- [70] Houston B, O'Neill IE, Mitchell MA, Goddard C. Purification and biological activity of a single charge isomer of pituitary-derived chicken growth hormone. J Endocrinol 1990;125:207–15.